
AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

- (Currently Amended) An absorbent article, preferably hygienic disposable absorbent article, comprising an agent able to convey a perception to the wearer, without the need to create the external condition perceived by the wearer.
- 2. (Currently Amended) An The article according to claim 1, wherein said article is a feminine hygienic disposable article, typically a panty liner, a sanitary napkin, an incontinent pad, an interlabial pad or a breast pad.
- 3. (Currently Amended) An <u>The</u> article according to claim 1, wherein the agent is a cooling agent able to stimulate thermo-receptors of the skin and/or mucosal surface of the wearer of the article, to convey a freshness sensation to the wearer, without the need to change the temperature on the skin and/or mucosal surface.
- 4. (Currently Amended) An The article according to claim 3 wherein the cooling agent is selected from the group consisting of ketals, carboxamides, cyclohexyl derivatives, cyclohexanol derivatives, camphor, borneol, eucalyptol, methyl salicylate, tea tree oil, eucalyptus oil, peppermint oil and mixtures thereof.
- 5. (Currently Amended) An The article according to claim 3, wherein the cooling agent is selected from the group consisting of:
 - a ketal according to the following formula:



in which R^1 represents a C_2 - C_6 -alkylene radical having at least 1, but not more than 3, hydroxyl group(s), preferably 1 hydroxyl group, and either R^2 and R^3 independently of one another represent C_1 - C_{10} -alkyl which is optionally substituted by about 1 to about 3

radicals selected from the group comprising hydroxyl, amino and halogen, C_5 - C_7 -cycloalkyl, preferably cyclohexyl, C_6 - C_{12} -aryl, preferably phenyl, with the proviso that the total of the C atoms of R^2 and R^3 is not less than 3, or R^2 and R^3 together represent an alkylene radical which, together with the carbon atom which carries the radicals R^2 and R^3 , forms a 5-7-membered ring, it being possible for this alkylene radical, in turn, to be substituted by C_1 - C_6 -aklyl groups, or mixtures thereof;

- or a carboxamide of the following formula:

(a)

wherein R', when taken separately, is hydrogen or an aliphatic radical containing up to about 25 carbon atoms; R" when taken separately is hydroxy, or an aliphatic radical containing up to about 25 carbon atoms, with the proviso that when R' is hydrogen, R" may also be an aryl radical of up to about 10 carbon atoms and selected from the group consisting of substituted phenyl, phenalkyl or substituted phenalkyl, naphthyl and substituted naphthyl, pyridyl; and R' and R", when taken together with the nitrogen atom to which they are attached, represent a cyclic or heterocyclic group of up to about 25 carbon atoms,

or (b)

$$R_1$$

$$R_2 \longrightarrow C^* \longrightarrow CONR'R"$$

$$R_3$$

wherein R' and R", when taken separately, are each hydrogen, C1-C5 alkyl or C1-C8 hydroxyalkyl and provide a total of no more than about 8 carbon atoms, with the proviso that when R' is hydrogen, R" may also be alkylcarboxyalkyl of up to about 6 carbon atoms; R' and R", when taken together, represent an alkylene group of up to 6 carbon atoms, the opposite ends of which group are attached to the amide nitrogen atom thereby to form a nitrogen heterocycle, the carbon chain of which may optionally be interrupted by oxygen; R1 is hydrogen or C1-C5 alkyl; and R2 and R3 are each C1-C5 alkyl; with the provisos that (i) R1, R2 and R3 together provide a total of at least about 5 carbon atoms, preferably from about 5-10 carbon atoms; and (ii) when R1 is hydrogen, R2 is C2 -C5 alkyl and R3 is C3-C5 alkyl and at least one of R2 and R3 is branched, preferably in an alpha or beta position relative to the carbon atom marked (*) in the formula, or a mixture thereof;

- or a cyclohexanol derivative according to the following general formulae:

wherein R represents a linear or branched alkyl group having about 1 to about 5 carbon atoms,

or (2)

wherein R¹ and R² are independently hydrogen, or a linear or branched alkyl group having about 1 to about 5 carbon atoms, or mixtures thereof;

- or a cyclohexyl derivative according to the following general formula

$$OR$$
 CH_2R_1

wherein R represents –H, a C_1 - C_5 linear or branched alkyl group, a C_1 - C_5 alkenyl group, a C_1 - C_5 alkoxy group or a C_1 - C_5 acyloxy group, R_1 represents –H, or a linear or branched alkyl group having from 1 to 5 carbon atoms, or mixtures thereof;

- or a mixture thereof.
- 6. (Currently Amended) An <u>The</u> article according to claim 3, wherein the cooling agent is a cyclohexyl derivative and more preferably is menthyl lactate.

- 7. (Currently Amended) An The article according to claim 3, which comprises menthol as a first cooling agent in combination with a second cooling agent, preferably selected from the group consisting of ketals, carboxamides, cyclohexyl derivatives other than menthol, cyclohexanol derivatives and mixtures thereof, preferably the second cooling agent is according to those of claim 5.
- 8. (Currently Amended) An The article according to claim 3, which comprises on at least a portion of the article from about 0.01 gm⁻² to about 300 gm⁻² of a cooling agent or a mixture thereof.
- 9. (Currently Amended) An <u>The</u> article according to any of the preceding claims 1 or 3, comprising a delivery system for containing and delivering the agent to at least a portion of the skin and/or mucosal surface of mammal wearing the article.
- 10. (Currently Amended) An <u>The</u> article according to claim 9, characterized in that wherein the delivery system is an emollient-containing composition comprising from about 0.1% to about 99.9%, by weight of the agent or mixture thereof, and from about 99.9% to about 0.1%, by weight of the emollient or mixture thereof.
- (Currently Amended) An The article according to claim 10, wherein the delivery system comprises an emollient typically selected from the group consisting of petroleum-based; fatty acids, particularly those having from about 8 to about 28 carbon atoms in their fatty chain; sucrose ester fatty acids; polyethylene glycol and derivatives thereof; fatty acid ester type; alkyl ethoxylate type; fatty acid ester ethoxylates; fatty alcohol type; polysiloxane type; propylene glycol and derivatives thereof; glycerine and derivatives thereof, including glyceride, acetoglycerides, and ethoxylated glycerides of C8-C28 fatty acids; spermaceti and other waxes; fatty alcohol ethers, particularly those having from about 8 to about 28 carbon atoms in their fatty chain, propoxylated fatty alcohols; other fatty esters of polyhydroxy alcohols; lanolin and its derivatives; kaolin and its derivatives; sorbitol and its derivative; trihydroxy stearin; ester derivatives and mixtures thereof.
- 12. (Currently Amended) An <u>The</u> article according to claim 9, wherein the delivery system comprises an alcohol or ester or acid derivative of following formulae or mixtures thereof

$$R_{6} - C - A = \begin{bmatrix} R_{4} \\ C \\ O \\ R_{1} \end{bmatrix} + \begin{bmatrix} R_{4} \\ C \\ O \\ R_{2} \end{bmatrix} + \begin{bmatrix} R_{4} \\ C \\ C \\ R_{2} \end{bmatrix} + \begin{bmatrix} R_{4} \\ R_{2} \end{bmatrix}$$

(I)

or

$$\begin{array}{c|c}
R_{8} - C - A \downarrow & R_{9} \\
C - A \downarrow & C \downarrow \\
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O$$

$$\begin{array}{c|c}
C = O & C = O
\end{array}$$

$$\begin{array}{c|c}
C = O & C = O$$

$$\begin{array}{c|c}
C = O &$$

wherein in (I) and (II) R_1 and each R_2 independently are an acyl group with from about 2 to about 22 carbon atoms, or an alkyl, alkenyl, arylalkyl, hydroxyalkyl group with from about 1 to about 24 carbon atoms or hydrogen, whereby at least one of R_1 and R_2 is such an acyl group, R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , and R_9 are independently an alkyl, alkenyl, arylalkyl, hydroxyalkyl, alkoxy groups of from about 1 to about 24 carbon atoms, hydroxy group or hydrogen group; R_{10} and R_{11} are independently an alkyl, alkenyl, arylalkyl, hydroxyalkyl, alkoxy groups of from about 2 to about 24 carbon atoms, hydroxy group or hydrogen group; A and B are independently a C_1 - C_6 linear or branched alkylene, alkyl, alkenylene, alkoxylene, alkoxyl, hydroxyalkylene, hydroxyalkyl groups; the values of x are independently from 0 to about 15; the values of y are independently 0 or 1,

or

wherein in (III) R_1 , R_2 and R_3 are independently an acyl, alkyl or alkenyl or hydroxyalkyl group with from about 1 to about 22 carbon atoms, and R_4 , R_5 , R_6 , R_7 and R_8 are independently selected from the group consisting of C_1 - C_{10} linear or branched alkyl, alkenyl, acyl, hydroxyalkyl or alkoxy groups, hydroxy, chloride, bromide, amine or hydrogen.

- 13. (Currently Amended) An The article according to claim 9, wherein the delivery system comprises triethyl citrate, acetyl tributyl citrate, triacetyl citrate, O acetyl triethyl citrate, polyethylene glycol and/or propylene glycol.
- 14. (Currently Amended) An The article according to any of the preceding claims 1 or 3, wherein the article comprises a wearer-facing surface, characterized in that wherein at least a portion of the wearer-facing surface comprises the agent.
- 15. (Currently Amended) An The article according to claim 3, wherein said article is a hygienic absorbent article comprising a topsheet as a wearer facing surface, a backsheet as a garment facing surface and an absorbent core sandwiched between the topsheet and backsheet, said backsheet preferably being a breathable backsheet.
- 16. (Currently Amended) An <u>The</u> absorbent article according to claim 15, wherein said breathable backsheet comprises at least one layer selected from an apertured polymeric film or a 2-dimensional planar apertured film.

- 17. (Currently Amended) An The absorbent article according to claim 16, wherein said layer is a 2-dimensional planar apertured layer, wherein said apertures have an average diameter of from about 150 micrometers to about 1 micrometers.
- 18. (Currently Amended) An <u>The</u> absorbent article according to claim 16, wherein said layer is an apertured polymeric film, wherein said apertures have an average diameter of from about 100 micrometers to about 500 micrometers.
- 19. (Currently Amended) An The absorbent article according to claim 16, wherein said breathable backsheet comprises at least two layers, a first layer comprising an apertured layer and a second layer comprising a fibrous layer.
- 20. (Cancelled)
- 21. (Currently Amended) An The article according to any of the preceding claims 1 or 3, preferably feminine hygienic disposable absorbent article, wherein at least a portion of the article is provided in a color that further contributes to the perception conveyed to the wearer of the article.